

## ADDENDUM INDEX

EXH.#	DOCUMENT DESCRIPTION	PAGE #
1	University of Colorado Law School, Intermountain Oil and Gas BMP Project Web Page, January 27, 2015.	1-3
2	National Association of Regional Councils Report – Local, Regional, and State Government Perspectives on Hydraulic Fracturing-Related Oil and Gas Development; Prepared By Samuel Gallaher, PhD Student at School of Public Affairs, University of Colorado Denver, Graduate Research Fellow at the Buechner Institute of Governance (excerpts), full report available at: <a href="http://narc.org/wp-content/uploads/Government-Perspectives-on-Oil-and-Gas-Development-Full-Report-2013-Gallaher.pdf">http://narc.org/wp-content/uploads/Government-Perspectives-on-Oil-and-Gas-Development-Full-Report-2013-Gallaher.pdf</a>	4-16
3	Energy Boomtown & Natural Gas: Implications for Marcellus Shale Local Governments & Rural Communities; NERCRD Rural Development Paper No. 43, 63 pp., Prepared by Jeffrey Jacquet, January 2009 (excerpts), full report available at: <a href="http://aese.psu.edu/nercrd/publications/rdp/rdp43">http://aese.psu.edu/nercrd/publications/rdp/rdp43</a>	17-22
4	Colorado Oil & Gas Association Rule 510 Statement, Prepared By Jamie L. Jost, Managing Shareholder at Jost & Shelton Energy Group, P.C., General Counsel for The Colorado Oil & Gas Association.	23-32
5	The Center for Science and Democracy at the Union of Concerned Scientists Report – Science, Democracy, and Fracking: A Guide for Community Residents and Policy Makers Facing Decisions Over Hydraulic Fracturing.	33-52
6	State of Colorado, Colorado Department of Public Health and Environment Letter Regarding Earth Guardians Request for Rulemaking, April 7, 2014.	53-54
7	Los Angeles Times Article: Message is mixed on Fracking, July 28, 2013.	55-58
8	Colorado Oil and Gas Conservation Commission, 2 CCR 404-1 – Statement of Basis, Specific Statutory Authority, and Purpose Re New Rules and Amendments to Current Rules (2008 Amendments) (excerpts), full copy available at: <a href="http://cogcc.state.co.us/">http://cogcc.state.co.us/</a>	59-65

9	New York State Department of Health Study: A Public Health Review of High Volume Hydraulic Fracturing for Shale Gas Development, December 17, 2014 (excerpts), full report available at: <a href="http://www.health.ny.gov/press/reports/docs/high_volume_hydraulic_fracturing.pdf">http://www.health.ny.gov/press/reports/docs/high_volume_hydraulic_fracturing.pdf</a>	66-83
10	Physicians, Scientists and Engineers for Health/Energy: Impediments to Public Health Research on Shale (Tight) Oil and Gas Development, May 2013.	84-85
11	National Public Radio State Impact: Lifelong Gag Order Imposed on Two Kids in Fracking Case, By Susan Phillips, August 1, 2013.	86-88
12	Pro Publica: EPA's Abandoned Wyoming Fracking Study One Retreat of Many, By Abraham Lustgarten, July 3, 2013.	89-92
13	Longmont Times Article: Most Oil, Gas Measures Die During Colorado Legislature's 2013 Session, By John Fryar, May 8, 2013.	93-95
14	National Public Radio Broadcast: Close Encounters With Gas Well Pollution; Host Broadcasters: Melissa Block and Robert Siegel, May 15, 2012	96-102
15	Health Impact Assessment for Battlement Mesa, Garfield County, Colorado, conducted by members of the faculty and staff of the Department of Environmental and Occupational Health, Colorado School of Public Health (CSPH), September 2010 (excerpts), full report available at: <a href="http://www.garfield-county.com/public-health/documents/1%20%20%20Complete%20HIA%20without%20Appendix%20D.pdf">http://www.garfield-county.com/public-health/documents/1%20%20%20Complete%20HIA%20without%20Appendix%20D.pdf</a>	103-122
16	U.S. Environmental Protection Agency News Release: EPA Releases Draft Findings of Pavillion, Wyoming Ground Water Investigation for Public Comment and Independent Scientific Review; EPA Contact Larry Jackson; December 8, 2011.	123-124
17	Pro Publica: EPA Finds Compound Used in Fracking in Wyoming Aquifer, By Abraham Lustgarten, November 10, 2011.	125-126
18	U.S. Environmental Protection Agency Draft Report Regarding Pavillion, Wyoming Groundwater Investigation for Public Comment and Independent Scientific Peer Review, December 8, 2011; Contact Person Richard Mylott, Public Affairs.	127-135

19	Bloomberg BNA: EPA Says Wyoming to Complete Investigation Of Possible Contamination Near Pavillon, Wyoming, By Alan Kovski, June 21.	136-140
20	U.S. Environmental Protection Agency News Release: EPA Initiates Hydraulic Fracturing Study: Agency Seeks Input From Science Advisory Board, March 18, 2010; Contact Person Enesta Jones.	141
21	Pro Publica: EPA Wants to Look at Full Lifecycle of Fracking in New Study, By Nicholas Kusnetz, February 9, 2011.	142-143
22	U.S. Environmental Protection Agency: Power Point Slides Regarding EPA Study of Hydraulic Fracturing and Drinking Water Resources.	144-161
23	Akron Beacon Journal: Article Regarding EPA Study on Fracking Threat to Water Will Take Years, By Bob Downing, January 18, 2013.	162-164
24	Physicians, Scientists and Engineers for Health/Energy: Working Paper – Toward an Understanding of the Environmental and Public Health Impacts of Shale Gas Development: An Analysis of the Peer-Reviewed Scientific Literature, 2009-2014, By Jake Hays and Seth B.C. Shonkoff, January 2015.	165-184
25	Pro Publica Surveys Some Recent Research on Potential Health Implications of Hydro Fracking: Drilling for Certainty - The Latest in Fracking Health Studies, By Naveena Sadasivam, March 5, 2014.	185-188
26	Environmental Health Perspectives, Volume 123, Number 1, January 2015: Proximity to Natural Gas Wells and Reported Health Status-Results of a Household Survey in Washington County, Pennsylvania, By P. M. Rabinowitz, I. B. Slizovskiy, V. Lamers, S. J. Trufan, T. R. Holford, J. D. Dziura, P. N. Peduzzi, M. J. Kane, J. S. Reif, T. R. Weiss, and M. H. Stowe.	189-194
27	Environmental Health Perspectives, Volume 122, Issue 4, April 2014: Birth Outcomes and Maternal Residential Proximity to Natural Gas Development in Rural Colorado, By L. M. McKenzie, R. Guo, R. Z. Witter, D. A. Savitz, L. S. Newman, and J. L. Adgate.	195-204
28	Physicians, Scientists and Engineers for Health/Energy Water Studies Summary: Surface and Groundwater Contamination Associated with Modern Natural Gas Development, October 2014.	205-206

29	University of Colorado Boulder: CU-Boulder Researchers Confirm Leaks From Front Range Oil and Gas Operations, May 7, 2014.	207-210
30	Colorado State University Report Regarding Characterizing Air Emissions from Natural Gas Drilling and Well Completion Operations, By Jeff Collett, Department of Atmospheric Science.	211-237
31	Rocky Mountain Mineral Law Foundation – Federal Onshore Oil & Gas Pooling & Unitization, Book 1; Mineral Law Series, Volume 2014, Number 4; Article on Pooling and Unitization: A History Perspective and an Introduction to Basic Vocabulary, By Bruce M. Kramer.	238-264
32	U.S. Energy Information Administration: North Dakota Aims to Reduce Natural Gas Flaring, Principal Contributors: Philip Budzik and Michael Ford, October 20, 2014.	265-266
33	National Center for Biotechnology Information Abstract: Impacts of Gas Drilling on Human and Animal Health, 2012.	267-268
34	Denver Business Journal: KC Fed: 50% of Energy Firms Planning Big Spending Cuts, Layoffs This Year, By Heather Draper, January 15, 2015.	269-270
35	The Scottish Government – News: Moratorium Called on Fracking, January 28, 2015.	271-273
36	Declaration of Ava Farouche (with maps).	274-278
37	Colo. Rev. Stat. §§ 34-60-102, 34-60-106.	279-288

*Environ Health Perspect*; DOI:10.1289/ehp.1306722

## Birth Outcomes and Maternal Residential Proximity to Natural Gas Development in Rural Colorado

Lisa M. McKenzie,<sup>1</sup> Ruixin Guo,<sup>2</sup> Roxana Z. Witter,<sup>1</sup> David A. Savitz,<sup>3</sup> Lee S. Newman,<sup>1</sup> and John L. Adgate<sup>1</sup>

Author Affiliations [open](#)

 [PDF Version \(258 KB\)](#)

Abstract

About This Article

Supplemental Material

**Background:** Birth defects are a leading cause of neonatal mortality. Natural gas development (NGD) emits several potential teratogens, and U.S. production of natural gas is expanding.

**Objectives:** We examined associations between maternal residential proximity to NGD and birth outcomes in a retrospective cohort study of 124,842 births between 1996 and 2009 in rural Colorado.

**Methods:** We calculated inverse distance weighted natural gas well counts within a 10-mile radius of maternal residence to estimate maternal exposure to NGD. Logistic regression, adjusted for maternal and infant covariates, was used to estimate associations with exposure tertiles for congenital heart defects (CHDs), neural tube defects (NTDs), oral clefts, preterm birth, and term low birth weight. The association with term birth weight was investigated using multiple linear regression.

**Results:** Prevalence of CHDs increased with exposure tertile, with an odds ratio (OR) of 1.3 for the highest tertile (95% CI: 1.2, 1.5); NTD prevalence was associated with the highest tertile of exposure (OR = 2.0; 95% CI: 1.0, 3.9, based on 59 cases), compared with the absence of any gas wells within a 10-mile radius. Exposure was negatively associated with preterm birth and positively associated with fetal growth, although the magnitude of association was small. No association was found between exposure and oral clefts.

**Conclusions:** In this large cohort, we observed an association between density and proximity of natural gas wells within a 10-mile radius of maternal residence and prevalence of CHDs and possibly NTDs. Greater specificity in exposure estimates is needed to further explore these associations.

### Introduction

Approximately 3.3% of U.S. live-born children have a major birth defect ([Centers for Disease Control and Prevention 2013](#); [Parker et al. 2010](#)); these defects account for 20% of infant deaths as well as 2.3% of premature death and disability ([McKenna et al. 2005](#)). Oral clefts, neural tube defects (NTDs), and congenital heart defects (CHD) are the most common classes of birth defects ([Parker et al. 2010](#)). These defects are thought to originate in the first trimester as a result of polygenic inherited disease or gene–environment interactions ([Brent 2004](#)). Suspected nongenetic risk factors for these birth defects include folate deficiency ([Wald and Sneddon 1991](#)), maternal smoking ([Honein et al. 2006](#)), alcohol abuse and solvent use ([Romitti et al. 2007](#)), and exposure to benzene ([Lupo et al. 2010b](#); [Wennborg et al. 2005](#)), toluene ([Bowen et al. 2009](#)), polycyclic aromatic hydrocarbons (PAHs) ([Ren et al. 2011](#)), and petroleum-based solvents, including aromatic hydrocarbons ([Chevrier et al. 1996](#)). Associations between air pollution [volatile organic compounds (VOCs), particulate matter (PM), and nitrogen dioxide (NO<sub>2</sub>)] and low birth weight and preterm birth have been reported ([Ballester et al. 2010](#); [Brauer et al. 2008](#); [Dadvand et al. 2013](#); [Ghosh et al. 2012](#); [Llop et al.](#)

2010). Many of these air pollutants are emitted during development and production of natural gas (referred to herein as NGD), and concerns have been raised that they may increase risk of adverse birth outcomes and other health effects (Colborn et al. 2011; McKenzie et al. 2012). Increased prevalence of low birth weight and small for gestational age and reduced APGAR scores were reported in infants born to mothers living near NGD in Pennsylvania (Hill 2013).

Technological advances in directional drilling and hydraulic fracturing have resulted in a global boom of drilling and production of natural gas reserves [U.S. Energy Information Administration (EIA) 2011a, 2011b; Vidas and Hugman 2008]. NGD is an industrial process resulting in potential worker and community exposure to multiple environmental stressors (Esswein et al. 2013; King 2012; Witter et al. 2013). Diesel-powered heavy equipment is used for worksite development as well as transporting large volumes of water, sand, and chemicals to sites and for waste removal (Witter et al. 2013). It is increasingly common for NGD to encroach on populated areas, potentially exposing more people to air and water emissions as well as to noise and community-level changes that may arise from industrialization [Colorado Oil and Gas Conservation Commission (COGCC) 2009]. Studies in Colorado, Texas, Wyoming, and Oklahoma have demonstrated that NGD results in emission of VOCs, NO<sub>2</sub>, sulfur dioxide (SO<sub>2</sub>), PM, and PAHs from either the well itself or from associated drilling processes or related infrastructure (i.e., drilling muds, hydraulic fracturing fluids, tanks containing waste water and liquid hydrocarbons, diesel engines, compressor stations, dehydrators, and pipelines) (CDPHE 2007; Frazier 2009; Kembell-Cook et al. 2010; Olaguer 2012; Walther 2011; Zielinska et al. 2011). Some of these pollutants, such as toluene, xylenes, and benzene, are suspected teratogens (Lupo et al. 2010b; Shepard 1995) or mutagens (Agency for Toxic Substances and Disease Registry 2007) and are known to cross the placenta (Bukowski 2001), raising the possibility of fetal exposure to these and other pollutants resulting from NGD. Currently, there are few studies on the effects of air pollution or NGD on birth outcomes.

In this analysis, we explored the association between maternal exposure to NGD and birth outcomes, using a data set with individual-level birth data and geocoded natural gas well locations. We conducted a retrospective cohort study to investigate the association between density and proximity of natural gas wells within a 10-mile radius of maternal residences in rural Colorado and three classes of birth defects, preterm birth, and fetal growth.

## Methods

**Study population.** We used information available in the publically accessible Colorado Oil and Gas Information System (COGIS) to build a geocoded data set with latitude, longitude, and year of development (1996–2009) for all gas wells in rural Colorado (COGIS 2011). Live birth data were obtained from the Colorado Vital Birth Statistics (CDPHE, Denver, CO). Geocoded maternal addresses at time of birth were linked to the well locations. Distance of each maternal residence from all existing (not abandoned) natural gas wells within a 10-mile radius was then computed using spherically adjusted straight line distances. We conducted our analysis on the final de-identified database containing maternal and birth outcome data described below and distance to all wells within the 10-mile radius. The Colorado Multiple Institutional Review Board reviewed and approved our study protocol. Informed consent was not required.

We restricted analysis to births occurring from 1996 through 2009 to focus our analysis on growth of unconventional NGD, characterized by use of hydraulic fracturing and/or directional drilling (King 2012), which expanded rapidly in Colorado beginning around 2000 (COGIS 2011). We also restricted our analysis to rural areas and towns with populations of < 50,000 (excluding the Denver metropolitan area, El Paso County, and the cities of Fort Collins, Boulder, Pueblo, Grand Junction, and Greeley) in 57 counties to reduce potential for exposure to other pollution sources, such as traffic, congestion, and industry. The final study area included locations with and without NGD. We conducted a retrospective study on the resulting cohort of 124,842 live births to explore associations between birth outcomes and exposure to NGD operations. We restricted eligibility to singleton births and excluded the small proportion (< 5%) of nonwhite births because there were too few to analyze separately.

**Birth outcomes.** Identified birth outcomes were a) oral cleft, including cleft lip with and without cleft palate as well as cleft palate [International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) code 749.xx] (National Center for Health Statistics 2011); b) NTD, including anencephalus, spina bifida without anencephaly, and encephalocele (ICD-9-CM 740.xx, 741.xx, and 742.0); c) CHD, including transposition of great vessels, tetralogy of Fallot, ventricular septal defect, endocardial cushion defect, pulmonary valve atresia and stenosis, tricuspid valve atresia and stenosis, Ebstein's anomaly, aortic valve stenosis, hypoplastic left heart syndrome, patent ductus arteriosus, coarctation of aorta, and pulmonary artery anomalies (codes 745.xx, 746.xx, 747.xx, excluding 746.9, 747.5); d) preterm birth (< 37 weeks completed gestation); e) term low birth weight ( $\geq$  37 weeks completed gestation and birth weight < 2,500 g); and f) term birth weight as a continuous measure. Births with an oral cleft, NTD, or CHD were excluded from preterm birth and term low birth weight analysis. Preterm births were excluded from term birth weight analysis. Oral cleft, CHD, and NTD cases in the Colorado Responds to Children with Special Needs (CRCSN) birth registry, obtained from

hospital records, the Newborn Genetics Screening Program, the Newborn Hearing Screening Program, laboratories, physicians, and genetic, developmental, and other specialty clinics (CRCSN 2011) were matched with Colorado live birth certificates. Cases are reflective of reporting as of 12 July 2012, were not necessarily confirmed by medical record review, and are subject to change as CRCSN ascertains diagnosis up to 3 years of child's age and/or supplements information by medical record review. We analyzed birth defects in three heterogeneous groups to increase statistical power. Data set information was not sufficient to distinguish between multiple and isolated birth anomalies or to identify chromosomal birth anomalies. In an exploratory analysis, we considered seven clinical diagnostic groupings of CHDs: a) conotruncal defects (tetralogy of Fallot and transposition of great vessels); b) endocardial cushion and mitral valve defects (EMD; endocardial cushion defect and hypoplastic left heart syndrome); c) pulmonary artery and valve defects (PAV; pulmonary valve atresia and stenosis and pulmonary artery anomalies); d) tricuspid valve defects (TVD; tricuspid valve atresia and stenosis and Ebstein's anomaly); e) aortic artery and valve defects (aortic valve stenosis and coarctation of aorta); f) ventricular septal defects (VSD); and g) patent ductus arteriosus in births > 2,500 g (Gilboa et al. 2005).

**Exposure assessment.** Distribution of the wells within a 10-mile radius of maternal residence shows 50% and 90% of wells to be within 2.3 and 7.7 miles of maternal residence, respectively. We used an inverse distance weighted (IDW) approach, commonly used to estimate individual air pollutant exposures from multiple fixed locations (Brauer et al. 1998; Ghosh et al. 2012), to estimate maternal exposure. Our IDW well count accounts for the number of wells within the 10-mile radius of the maternal residence, as well as distance of each well from the maternal residence, giving greater weight to wells closest to the maternal residence. For example, an IDW well count of 125 wells/mile could be computed from 125 wells each located 1 mile from the maternal residence or 25 wells each located 0.2 miles from the maternal residence. We calculated the IDW well count of all existing natural gas wells in the birth year within a 10-mile radius of each maternal residence as a continuous exposure metric:

$$\text{IDW well count} = \sum_{i=1}^n \frac{1}{d_i} \quad [1]$$

where IDW well count is the IDW count of existing wells within a 10-mile radius of maternal residence in the birth year;  $d_i$  is the distance of the  $i$ th individual well from maternal residence; and  $n$  is the number of existing wells within a 10-mile radius of maternal residence in the birth year.

The IDW well count was calculated for each maternal residence with  $\geq 1$  gas wells within 10 miles. The final distribution then was divided into tertiles (low, medium, and high) for subsequent logistic and linear regression analysis. Each tertile was compared with the referent group (no natural gas wells within 10 miles, IDW well count = 0).

**Statistical analysis.** We used logistic regressions to study associations between each dichotomous outcome and IDW exposure group. We also considered term birth weight as a continuous outcome using multiple linear regression. First, we estimated the crude odds ratio (OR) associated with IDW exposure tertile for each binary outcome, followed by a Cochran–Armitage test to evaluate linear trends in binomial proportions with increasing IDW exposure (none, low, medium, and high). We further investigated associations by adjusting for potential confounders, as well as infant and maternal covariates selected based on both *a priori* knowledge and empirical consideration of their association with exposure and an outcome. Specifically, covariates in our analysis of all outcomes except outcomes with very few events (i.e., NTDs, conotruncal defects, EMDs, and TVDs) included maternal age, education (< 12, 12, 13–15,  $\geq 16$  years), tobacco use (smoker, nonsmoker), ethnicity (Hispanic, non-Hispanic white), and alcohol use (yes, no), as well as parity at time of pregnancy (0, 1, 2, > 2) and infant sex. Gestational age was also included in the analysis of term birth weight. Elevation of maternal residence also was considered in the analysis because most wells are < 7,000 feet, and elevation has been associated with both preterm birth and low birth weight (Niermeyer et al. 2009). For 272 births where elevation of maternal residence was missing, elevation was imputed using mean elevation for maternal ZIP code. For outcomes with very few events, only elevation was included in the multiple logistic modeling to avoid unstable estimates. The ORs and their 95% CIs were used to approximate relative risks for each outcome associated with IDW count exposure tertile (low, medium, and high) compared with no wells within 10 miles, which is reasonable because of the rarity of the outcomes. We considered the statistical significance of the association, as well as the trend, in evaluating results, at an alpha of 0.05. We evaluated the confounding potential of the 1998 introduction of folic acid fortification on the birth defect outcomes and found only a decrease in NTD prevalence after 1998 (see Supplemental Material, Table S1).

In a sensitivity analyses, we explored reducing exposure to 2- and 5-mile buffers around the maternal residence, as well as restricting the cohort to births occurring between 2000 and 2009 to exclude births before the expansion of NGD. We report estimated associations with 95% CIs. All statistical analyses were conducted using SAS® software version 9.3 (SAS Institute Inc., Cary, NC).

## Results

Births were approximately evenly divided between exposed and unexposed groups (0 wells in a 10-mile radius versus  $\geq 1$  well in a 10-mile radius) (Table 1). Estimated exposure, represented by IDW well counts, tended to be higher for births to mothers with residence addresses at lower elevations ( $< 6,000$  feet), and among nonsmoking and Hispanic mothers (Table 1).

Table 1 – Study population characteristics for unexposed and exposed subjects from rural Colorado 1996–2009.

[View Table \(HTML Version\)](#)  
[View larger image \(TIF File\)](#)

Both crude and adjusted estimates indicate a monotonic increase in the prevalence of CHDs with increasing exposure to NGD, as represented by IDW well counts (Table 2). Births to mothers in the most exposed tertile ( $> 125$  wells/mile) had a 30% greater prevalence of CHDs (95% CI: 1.2, 1.5) than births to mothers with no wells within a 10-mile radius of their residence.

Table 2 – Association between inverse distance weighted well count within 10-mile radius of maternal residence and CHDs, NTDs, and oral clefts.

[View Table \(HTML Version\)](#)  
[View larger image \(TIF File\)](#)

Prevalence of NTDs was positively associated with only the third exposure tertile, based on crude and estimated adjusted ORs for elevation (Table 2). Births in the highest tertile ( $> 125$  wells/mile) were 2.0 (95% CI: 1.0, 3.9) times more likely to have a NTD than those with no wells within a 10-mile radius, based on 59 available cases. We observed no statistically significant associations between oral clefts and NGD, based on trend analysis across categorical IDW well count exposure (Table 2).

Both crude and adjusted estimates for preterm birth suggest a slight ( $< 10\%$ ) decreased risk of preterm birth with increasing exposure to NGD (Table 3). Crude term low birth weight measures suggested decreased risk of term low birth weight with increasing exposure to NGD. A weak nonlinear trend remained after adjusting for elevation and other covariates. This association is consistent with the multiple linear regression results for continuous term birth weight, in which mean birth weights were 5–24 g greater in the higher IDW well count exposure tertiles than the referent group.

Table 3 – Association between inverse distance weighted well count within 10-mile radius of maternal residence and preterm birth and term low birth weight.

[View Table \(HTML Version\)](#)  
[View larger image \(TIF File\)](#)

We observed a monotonic increase in the prevalence of NTDs with increasing exposure to NGD in our sensitivity analyses using 2- and 5-mile exposure radii as well as some attenuation in decreased risk for preterm birth and term low birth weight (see Supplemental Material, Tables S2–7). Restricting births to 2000 through 2009, the period of most intense NGD in Colorado, attenuated the positive association between NTDs in the highest tertile and did not alter observed relationships for other birth outcomes (see Supplemental Material, Tables S2–S7).

Exploratory analysis of CHDs by clinical diagnostic groups indicates increased prevalence of PAV defects by 60% (95% CI: 1.1, 2.2), VSDs by 50% (95% CI: 1.1, 2.1), and TVDs by 400% (95% CI: 1.3, 13) in the most exposed tertile compared with those with no wells within a 10-mile radius (Table 4).

Table 4 – Association between inverse distance weighted well count within 10-mile radius of maternal residence and CHD diagnostic groups.

[View Table \(HTML Version\)](#)  
[View larger image \(TIF File\)](#)

## Discussion

We found positive associations between density and proximity of natural gas wells within a 10-mile radius of maternal residence and birth prevalence of CHDs and possibly NTDs. Prevalence of CHDs increased monotonically from the lowest to highest exposure tertile, although even in the highest tertile the magnitude of the association was modest. Prevalence of NTDs was elevated only in the highest tertile of exposure. We also observed small negative associations between density and proximity of natural gas wells within a 10-mile radius of maternal residence and preterm birth and term low birth weight, and a small positive association with mean birth weight. We found no indication of an association between density and proximity of natural gas wells within a 10-mile radius of maternal residence and oral cleft prevalence.

Nongenetic risk factors for CHDs and NTDs possibly attributable to NGD include maternal exposure to benzene ([Lupo et al. 2010b](#); [Wennborg et al. 2005](#)), PAHs ([Ren et al. 2011](#)), solvents ([Brender et al. 2002](#); [Chevrier et al. 1996](#); [Desrosiers et al. 2012](#); [McMartin et al. 1998](#)), and air pollutants (NO<sub>2</sub>, SO<sub>2</sub>, PM) ([Vrijheid et al. 2011](#)). NGD emits multiple air pollutants, including benzene and toluene, during the “well completion” phase (when gas and water flow back to the surface after hydraulic fracturing) as well as from related infrastructure ([CDPHE 2009a, 2009b](#); [Garfield County Public Health Department 2009](#); [Gilman et al. 2013](#); [McKenzie et al. 2012](#); [Pétron et al. 2012](#)). Ambient benzene levels in areas with active NGD in Northeast Colorado ranged from 0.03 to 6 parts per billion by volume (ppbv) ([CDPHE 2012](#); [Gilman et al. 2013](#); [Pétron et al. 2012](#)). Furthermore, 24-hr average ambient air benzene levels near active well development sites in western Colorado ranged from 0.03 to 22 ppbv ([McKenzie et al. 2012](#)).

Two previous case-control studies have reported associations between maternal exposure to benzene and birth prevalence of NTDs and/or CHDs ([Lupo et al. 2010b](#); [Wennborg et al. 2005](#)). The study by [Lupo et al. \(2010b\)](#) of 4,531 births in Texas found that mothers living in census tracts with the highest ambient benzene levels (0.9–2.33 ppbv) were 2.3 times more likely to have offspring with spina bifida than mothers living in census tracts with the lowest ambient benzene levels (95% CI: 1.22, 4.33). An occupational study of Swedish laboratory employees found a significant association between exposure to occupational levels of benzene in the critical window between conception, organogenesis, and neural crest formation and neural crest malformations ([Wennborg et al. 2005](#)). Children born to 298 mothers exposed to benzene had 5.3 times greater prevalence of neural crest malformations than children born to mothers not exposed to benzene (95% CI: 1.4, 21.1). Other studies of maternal exposures to organic solvents, some of which contain benzene, have reported associations between maternal occupational exposure to organic solvents and major birth defects ([Brender et al. 2002](#); [Desrosiers et al. 2012](#); [McMartin et al. 1998](#)). Although exposure to benzene is a plausible explanation for the observed associations, further research is needed to examine whether these associations are replicated and whether benzene specifically explains these associations.

Air pollutants emitted from diesel engines used extensively in NGD also may be associated with CHDs and/or NTDs. Trucks with diesel engines are used to transport supplies, water, and waste to and from gas wells, with 40 to 280 truck trips per day per well pad during development ([Witter et al. 2013](#)). Generators equipped with diesel engines are used in both drilling wells and hydraulic fracturing. Air pollutants in diesel exhaust include NO<sub>2</sub>, SO<sub>2</sub>, PM, and PAHs. A meta-analysis of four studies suggested associations of maternal NO<sub>2</sub> and SO<sub>2</sub> exposures with coarctation of the aorta and tetralogy of Fallot, and of maternal PM<sub>10</sub> exposure with arterial septal defects ([Vrijheid et al. 2011](#)). Two case-control studies in China reported positive associations between PAH concentrations in maternal blood and the placenta and NTDs ([Li et al. 2011](#); [Naufal et al. 2010](#)). Several CHDs were associated with traffic related carbon monoxide and ozone pollution in a case control study of births from 1987 to 1993 in Southern California ([Ritz et al. 2002](#)).

The small negative associations with term low birth weight and preterm birth in our study population were unexpected given that other studies have reported positive associations between these outcomes and urban air pollution ([Ballester et al. 2010](#); [Brauer et al. 2008](#); [Dadvand et al. 2013](#); [Ghosh et al. 2012](#); [Llop et al. 2010](#)) and proximity to natural gas wells ([Hill 2013](#)). It is possible that rural air quality near natural gas wells in Colorado is not as compromised as urban air quality in these studies, and exposure represented as IDW well count may not adequately represent air quality. In addition, the power of our large cohort increases the likelihood of false positive results for small associations close to the null. Although associations were consistent across measures of birth weight (i.e., reduced risk of term low birth weight and increase in mean birth weight), they attenuated toward the null in sensitivity analysis for 2- and 5-mile radii (see Supplemental Material, Tables S6–S7). If causal, stronger associations would be expected with more stringent exposure definitions. Our incomplete ability to adjust for socioeconomic status, health, nutrition, prenatal care, and pregnancy complications likely accounts for these unexpected findings.

This study has several limitations inherent in the nature of the available data. Not all birth defects were confirmed by medical record review. Also, birth defects are most likely undercounted, because stillbirths, terminated pregnancies, and later-life diagnoses (after 3 years of age) are not included. Birth weight and gestational age were obtained from birth

certificates, which are generally accurate for birth weight and useful but less accurate for gestational age (DiGiuseppe et al. 2002). Data on covariates were obtained from birth certificates and were limited to basic demographic, education, and behavioral information available in the vital records. Distribution of covariates among exposure tertiles and the unexposed group was similar; nevertheless, our incomplete ability to adjust for socioeconomic status, health, nutrition, prenatal care, and pregnancy complications may have resulted in residual confounding. In addition, low event outcomes (e.g., NTDs) were adjusted only for elevation. The data set did not contain information on maternal folate consumption and genetic anomalies, both independent predictors of our outcomes, which may have confounded these results. We did observe a large decrease in the prevalence of NTDs after the introduction of folic acid in 1998, and small increases in the prevalence of CHDs and oral clefts, although none of the estimates are statistically significant (see Supplemental Material, Table S1). Further study is needed to determine whether unaccounted folate confounding is attenuating our results toward the null. There is no evidence indicating genetic anomalies would differ by IDW well count around maternal residence.

Because of the rarity of specific birth defects in the study population, birth defects were aggregated into three general groups. This limited our study in that associations with specific birth defects may have been obscured. An exploratory analysis of CHDs by clinical diagnostic groups indicates increased prevalence of specific diagnostic groups (i.e., PAV, VSD, and TVD) compared with aggregated CHDs (Table 4).

Another limitation of this study is the lack of temporal and spatial specificity of the exposure assessment. Because we did not have maternal residential history, we assumed that maternal address at time of delivery was the same as maternal address during the first trimester of pregnancy—the critical time period for formation of birth defects. Studies in Georgia and Texas estimate that 22–30% of mothers move residence during their pregnancy, and most mothers move within their locality (Lupo et al. 2010a; Miller et al. 2010), potentially introducing some exposure misclassification for the early pregnancy period of interest. However, these studies found little difference in mobility between cases and controls (Lupo et al. 2010a; Miller et al. 2010), and maternal mobility did not significantly influence the assessment of benzene exposure (Lupo et al. 2010a). We were able to determine only whether a well existed within the calendar year of birth (e.g., 2003) and did not have sufficient data to determine if a well existed within the first trimester of the pregnancy. Therefore, some nondifferential exposure misclassification is likely and the overall effect of this is unknown.

Similarly, we had consistent information only on existence of a well in the birth year. Lack of information on natural gas well activity levels, such as whether or not wells were producing or undergoing development, may have resulted in exposure misclassification. Actual exposure to natural gas–related pollutants likely varies by intensity of development activities. Lack of temporal and spatial specificity of the exposure assessment would most likely have tended to weaken associations (Ritz et al. 2007; Ritz and Wilhelm 2008). To address spatial and temporal variability, additional air pollution measurements and modeling will be needed to improve exposure estimates at specific locations. Last, information on the mother's activities away from her residence, such as work and recreation, as well as proximity of these activities to NGD was not available and may have led to further exposure misclassification and residual confounding.

## Conclusion

This study suggests a positive association between greater density and proximity of natural gas wells within a 10-mile radius of maternal residence and greater prevalence of CHDs and possibly NTDs, but not oral clefts, preterm birth, or reduced fetal growth. Further studies incorporating information on specific activities and production levels near homes over the course of pregnancy would improve exposure assessments and provide more refined effect estimates. Recent data indicate that exposure to NGD activities is increasingly common. The COGCC estimates that 26% of the > 47,000 oil and gas wells in Colorado are located within 150–1,000 feet of a home or other type of building intended for human occupancy (COGCC 2012). Taken together, our results and current trends in NGD underscore the importance of conducting more comprehensive and rigorous research on the potential health effects of NGD.

## References

Agency for Toxic Substances and Disease Registry. 2007. Toxicological Profile for Benzene. Available: <http://www.atsdr.cdc.gov/ToxProfiles/TP.asp?id=40&tid=14> [accessed 22 May 2013].

Ballester F, Estarlich M, Iniguez C, Llop S, Ramon R, Esplugues A, et al. 2010. Air pollution exposure during pregnancy and reduced birth size: a prospective birth cohort study in Valencia, Spain. *Environ Health* 9:6; doi: [10.1186/1476-069X-9-6](https://doi.org/10.1186/1476-069X-9-6).

- Bowen S, Irtenkauf S, Hannigna J, Stefanski A. 2009. Alterations in rat morphology following abuse patterns of toluene exposure. *Reprod Toxicol* 27:161–169.
- Brauer M, Lencar C, Tamburic L, Koehoorn M, Demers P, Karr C. 2008. A cohort study of traffic-related air pollution impacts on birth outcomes. *Environ Sci Technol* 116:680–686.
- Brender JD, Suarez L, Hendricks KA, Baetz RA, Larsen R. 2002. Parental occupation and neural tube defect-affected pregnancies among Mexican Americans. *J Occup Environ Med* 44:650–656.
- Brent RL. 2004. Environmental causes of human congenital malformations: the pediatrician's role in dealing with these complex clinical problems caused by a multiplicity of environmental and genetic factors. *Pediatrics* 113:957–968.
- Bukowski JA. 2001. Review of the epidemiological evidence relating toluene to reproductive outcomes. *Regul Toxicol and Pharmacol* 33:147–156.
- CDPHE (Colorado Department of Public Health and Environment). 2007. Garfield County Air Toxics Inhalation: Screening Level Human Health Risk Assessment: Inhalation Of Volatile Organic Compounds Measured In Rural, Urban, and Oil & Gas Areas In Air Monitoring Study (June 2005–May 2007). Available: <http://www.garfield-county.com/public-health/documents/Working%20Draft%20CDPHE%20Screening%20Level%20Risk%20Air%20Toxics%20Assessment%2012%2020%2007.pdf> [accessed 22 May 2013].
- CDPHE (Colorado Department of Public Health and Environment). 2009a. Garfield County Emissions Inventory. Available: [http://www.garfield-county.com/air-quality/documents/airquality/Garfield\\_County\\_Emissions\\_Inventory-2009.pdf](http://www.garfield-county.com/air-quality/documents/airquality/Garfield_County_Emissions_Inventory-2009.pdf) [accessed 22 May 2013].
- CDPHE (Colorado Department of Public Health and Environment). 2009b. State of Colorado Technical Support Document for Recommended 8-Hour Ozone Designations. Available: <http://www.colorado.gov/cs/Satellite?blobcol=urldata&blobheadername1=Content-Disposition&blobheadername2=Content-Type&blobheadervalue1=inline%3B+filename%3D%22Recommended+8+Hour+Ozone+Area+Designation.s.pdf%22&blobheadervalue2=application%2Fpdf&blobkey=id&blobtable=MungoBlobs&blobwhere=1251808873080&ssbinary=true> [accessed 22 May 2013].
- CDPHE (Colorado Department of Public Health and Environment). 2012. Air Emissions Case Study Related to Oil and Gas Development in Erie, Colorado. Available: [http://www.colorado.gov/airquality/tech\\_doc\\_repository.aspx](http://www.colorado.gov/airquality/tech_doc_repository.aspx) [accessed 22 May 2013].
- Centers for Disease Control and Prevention. 2013. Birth Defects. Available: <http://www.cdc.gov/ncbddd/birthdefects/index.html> [accessed 22 May 2013].
- Chevrier C, Dananche B, Bahuau M, Nelva A, Herman C, Francannet C, et al. 1996. Occupational exposure to organic solvent mixtures during pregnancy and the risk of non-syndromic oral clefts. *Occup Environ Med* 63:617–623.
- COGCC (Colorado Oil and Gas Conservation Commission). 2009. Statement of Basis, Specific Statutory Authority, and Purpose: New Rules and Amendments to Current Rules of the Colorado Oil and Gas Conservation Commission, 2 ccr 404-1. Available: <http://cogcc.state.co.us/> [accessed 22 May 2013].
- COGCC (Colorado Oil and Gas Conservation Commission). 2012. Staff Report. Colorado Department of Natural Resources. Available: <http://cogcc.state.co.us/> [accessed 22 May 2013].
- COGIS (Colorado Oil and Gas Information System). 2011. Well Production Database. Vol. 2011. Available: <http://cogcc.state.co.us/> [accessed 22 May 2013].
- Colborn T, Kwiatkowski C, Schultz K, Bachran M. 2011. Natural gas operations from a public health perspective. *Hum Ecol Risk Asses* 17:1039–1056.
- CRCSN (Colorado Responds to Children with Special Needs). 2011. Birth Defects Dataset Details. Available: <http://www.cdphe.state.co.us/cohid/crcsndata.html> [accessed 6 October 2011].
- Dadvand P, Parker JD, Bell ML, Bonzini M, Brauer M, Darrow L, et al. 2013. Maternal exposure to particulate air pollution and term birth weight: a multi-country evaluation of effect and heterogeneity. *Environ Health Perspect* 121:367–373; doi: [10.1289/ehp.1205575](https://doi.org/10.1289/ehp.1205575).
- Desrosiers TA, Lawson CC, Meyer RE, Richardson DB, Daniels JL, Waters MA, et al. 2012. Maternal occupational exposure to organic solvents during early pregnancy and risks of neural tube defects and orofacial clefts. *Occup Environ Med* 69:493–499.