

D. Reporting templates

Based on the need to now report data not only for FC citywide but also separately for each of the four zones (NW, NE, SE, SW; broken down using College and Drake) the city has been divided into for the purpose of WNV surveillance and mosquito control, CSU had to generate a new reporting template. This new reporting template is tailored to provide data relevant to the Level III and IV entomological triggers (see below) for control measures in the “City of Fort Collins Program Response Guidelines to Mosquito Borne Arboviral Activity (March 2013 edition)”.

Level III

- Vector index > 0.75 and increasing
- Mosquito infection rates of > 3.0 per thousand (0.3%) and increasing

Level IV

- Vector index > 0.75.
- Sustained mosquito infection rates of > 5.0 per thousand (0.5%)

The new reporting format comprises a set of 6 tables to address the current week (1a, 2a, 3a) and to provide seasonal and historical context (1b, 2b, 3b) (see full table formats on following pages)

- Table 1a. Vector Index for current week
- Table 1b. Vector Index for All *Culex* by week from June-August
- Table 2a. Vector abundance for current week
- Table 2b. Vector abundance for All *Culex* by week from June-August
- Table 3a. WNV infection rate per 1,000 females for current week
- Table 3b. WNV infection rate per 1,000 females for All *Culex* by week from June-August

Table 1a. Vector Index for current week

Week: 34	Mean abundance of females per trap night ¹		Estimate for proportion of females infected with WNV ²		Vector Index		
	Cx. <i>pipiens</i>	Cx. <i>tarsalis</i>	Cx. <i>pipiens</i>	Cx. <i>tarsalis</i>	Cx. <i>pipiens</i> ³	Cx. <i>tarsalis</i> ⁴	All <i>Culex</i> ⁵
FC – Zone NW	5.22	24.44	0	0	0	0	0
FC – Zone NE	22.60	83.20	0.00342	0.00120	0.077	0.100	0.177
FC – Zone SE	12.80	42.87	0.01207	0.00449	0.154	0.192	0.346
FC – Zone SW	2.38	7.25	0	0	0	0	0
FC – Citywide	11.52	41.74	0.00651	0.00228	0.075	0.095	0.170
LV	9.94	17.39	0	0.0158	0	0.275	0.275

¹From Table 2a (CDC light trap catches only).²Derived from the data presented in Table 3a for estimated infection rate per 1,000 females (CDC light trap and gravid trap catches combined).³Vector Index for Cx. *pipiens* = (Mean abundance of Cx. *pipiens* females per trap night) x (Estimate for proportion of all Cx. *pipiens* females infected with WNV).⁴Vector Index for Cx. *tarsalis* = (Mean abundance of Cx. *tarsalis* females per trap night) x (Estimate for proportion of all Cx. *tarsalis* females infected with WNV).⁵Vector Index for All *Culex* = (Vector Index for Cx. *pipiens*) + (Vector Index for Cx. *tarsalis*).**Table 1b. Vector Index for All *Culex* by week from June-August**

Week	FC – Zone NW		FC – Zone NE		FC – Zone SE		FC – Zone SW		FC – Citywide		LV
	Current year	Historical average ¹	Current year	Historical average ¹							
23	0	0.018	0	0.020	0	0.017	0	0.014	0	0.021	0
24	0	0	0	0	0	0.014	0	0	0	0.004	0
25	0	0	0	0.013	0	0	0	0	0	0.003	0
26	0	0	0	0	0.022	0	0.014	0	0.010	0	
27	0	0	0	0.034	0	0.011	0	0	0	0.012	0
28	0	0	0	0.037	0	0.127	0	0.012	0	0.051	0
29	0	0.126	0	0.164	0	0.355	0	0.041	0	0.193	0
30	0	0.123	0	0.168	0.203	0.304	0	0.102	0.070	0.191	0.118
31	NA	0.224	0.412	0.074	0.570	0.152	NA	0.072	NA	0.131	0.095
32	0.222	0.162	0.302	0.311	0.634	0.365	0.020	0.107	0.352	0.259	0.208
33	0.018	0.218	0.159	0.428	0.270	0.272	0.116	0.051	0.156	0.255	0.128
34	0	0.174	0.177	0.163	0.346	0.313	0	0.074	0.170	0.192	0.275
35		0.062		0.126		0.192		0.198		0.149	

¹2006-2013 (2003-2005 were excluded due to changes in trap locations from 2006 onwards).

Table 2a. Vector abundance for current week (CDC light trap catches only)

Week: 34	Total number females collected			Number CDC light trap nights	Mean abundance of females per CDC light trap night		
	Cx. <i>pipiens</i>	Cx. <i>tarsalis</i>	All <i>Culex</i>		Cx. <i>pipiens</i> ¹	Cx. <i>tarsalis</i> ²	All <i>Culex</i> ³
FC – Zone NW	47	220	267	9	5.22	24.44	29.67
FC – Zone NE	226	832	1058	10	22.60	83.20	105.80
FC – Zone SE	192	643	835	15	12.80	42.87	55.67
FC – Zone SW	19	58	77	8	2.38	7.25	9.62
FC – Citywide	484	1753	2237	42	11.52	41.74	53.26
LV	358	626	984	36	9.94	17.39	27.33

¹Mean abundance of *Cx. pipiens* females per CDC light trap night = (Total number *Cx. pipiens* females collected) / (Number CDC light trap nights).

²Mean abundance of *Cx. tarsalis* females per CDC light trap night = (Total number *Cx. tarsalis* females collected) / (Number CDC light trap nights).

³Mean abundance of All *Culex* females per CDC light trap night = (Total number All *Culex* females collected) / (Number CDC light trap nights).

Table 2b. Vector abundance for All *Culex* by week from June-August

Week	FC – Zone NW		FC – Zone NE		FC – Zone SE		FC – Zone SW		FC – Citywide		LV
	Current year	Historical average ¹	Current year	Historical average ¹							
23	3.22	2.02	1.10	5.86	2.27	8.59	1.00	0.22	1.93	5.21	3.11
24	1.33	2.31	0.90	8.73	1.20	12.88	0.67	0.63	1.05	7.07	2.00
25	3.44	6.77	6.90	18.15	6.60	23.08	0.44	2.13	4.72	14.20	6.03
26	2.44	12.68	11.50	38.60	16.07	35.39	2.22	6.27	9.26	25.01	9.84
27	22.56	31.85	61.90	69.76	58.27	40.90	12.89	12.48	42.14	42.47	54.69
28	39.22	49.17	116.20	80.96	74.33	66.50	14.56	12.97	64.21	54.78	42.05
29	27.00	58.82	57.20	87.35	36.33	73.48	6.44	21.88	32.98	63.27	30.59
30	94.11	42.96	236.40	106.32	213.67	61.07	52.89	12.08	160.28	58.46	75.94
31	NA	45.53	263.90	74.63	191.23	43.18	NA	12.95	NA	45.38	83.86
32	27.56	43.06	165.90	55.75	198.20	38.77	26.67	13.12	119.07	38.66	51.59
33	38.33	25.43	69.40	49.56	90.27	28.43	26.56	6.90	61.21	28.28	37.86
34	29.67	18.62	105.80	38.14	55.67	23.15	9.62	6.11	53.26	22.45	27.33
35		8.27		18.56		9.84		3.65		10.36	

¹2006-2013 (2003-2005 were excluded due to changes in trap locations from 2006 onwards).

Table 3a. WNV infection rate per 1,000 females for current week (CDC light trap and gravid trap catches combined)

Week: 34	Total number individuals examined			Total number pools examined			Total number WNV-infected pools			Estimate for WNV infection rate per 1,000 females ¹		
	Cx. <i>pipiens</i>	Cx. <i>tarsalis</i>	All <i>Culex</i>	Cx. <i>pipiens</i>	Cx. <i>tarsalis</i>	All <i>Culex</i>	Cx. <i>pipiens</i>	Cx. <i>tarsalis</i>	All <i>Culex</i>	Cx. <i>pipiens</i>	Cx. <i>tarsalis</i>	All <i>Culex</i>
FC – Zone NW	73	220	293	10	10	20	0	0	0	0	0	0
FC – Zone NE	291	833	1124	16	22	38	1	1	2	3.42	1.20	1.81
FC – Zone SE	423	696	1119	22	23	45	4	3	7	12.07	4.49	7.11
FC – Zone SW	85	59	144	8	9	17	0	0	0	0	0	0
FC – Citywide	872	1808	2680	56	64	120	5	4	9	6.51	2.28	3.59
LV	12	154	166	6	7	13	0	2	2	0	15.80	14.54

¹Maximum likelihood estimate (MLE) for WNV infection rate per 1,000 females calculated using the CDC PooledInfRate 4.0 plug-in for Excel.

Table 3b. WNV infection rate per 1,000 females for All *Culex* by week from June-August

Week	FC – Zone NW		FC – Zone NE		FC – Zone SE		FC – Zone SW		FC – Citywide		LV
	Current year	Historical average ¹	Current year	Historical average ¹							
23	0	9.00	0	4.62	0	2.25	0	53.30	0	4.19	0
24	0	0	0	0	0	1.09	0	0	0	0.60	0
25	0	0	0	0.74	0	0	0	0	0	0.25	0
26	0	0	0	0	0	0.61	0	2.30	0	0.41	0
27	0	0	0	0.48	0	0.25	0	0	0	0.28	0
28	0	0	0	0.56	0	1.92	0	0.90	0	0.98	0
29	0	2.02	0	1.77	0	4.59	0	1.84	0	2.91	0
30	0	3.03	0	1.58	0.88	5.22	0	8.74	0.41	3.37	1.56
31	NA	5.06	1.52	1.08	2.88	3.69	NA	5.46	NA	2.95	1.14
32	7.87	4.07	2.18	5.74	3.32	9.70	3.20	8.23	3.18	6.83	4.72
33	1.64	8.85	2.54	8.26	3.05	11.00	2.41	7.68	2.65	9.24	3.95
34	0	9.82	1.81	4.49	7.11	12.13	0	12.41	3.59	8.68	14.54
35		7.66		6.80		20.00		53.69		14.33	

¹2006-2013 (2003-2005 were excluded due to changes in trap locations from 2006 onwards).